

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES		MALE	BOTH SEXES		MALE									
All causes															
2000.....	872.0	1,042.5	739.1	852.1	1,018.2	722.2	971.8	1,174.5	819.7	1,129.9	1,377.8	947.9			
1999.....	881.9	1,061.8	743.6	860.7	1,035.8	725.7	990.8	1,206.7	831.0	1,147.1	1,412.5	955.0			
I. Certain infectious and parasitic diseases (A00-B99)															
2000.....	21.4	26.3	17.0	17.5	21.3	14.0	43.9	56.5	33.6	55.7	73.0	41.8			
1999.....	22.1	27.4	17.4	17.9	22.0	14.3	46.3	60.1	35.0	57.7	76.3	42.9			
Intestinal infectious diseases (A00-A09)															
2000.....	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.3	0.3	0.4	0.4	0.3			
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3			
Cholera (A00)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*			
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			
Other intestinal infectious diseases (A01-A08)															
2000.....	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.3	0.3	0.3	0.4	0.3			
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.2	0.2	0.2	0.3	0.3	0.3			
Typhoid fever (A01.0)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*			
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)															
2000.....	0.0	*	*	0.0	*	*	*	*	*	*	*	*			
1999.....	0.0	0.0	*	0.0	*	*	*	*	*	*	*	*			
Shigellosis (A03)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*			
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Other bacterial food-borne intoxications (A05)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Amebiasis (A06)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Intestinal infections due to other specified organisms (A04,A07-A08)															
2000.....	0.5	0.5	0.4	0.5	0.5	0.5	0.3	0.3	0.3	0.3	0.3	0.4	0.3		
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.2	0.2	0.2	0.3	0.3	0.3	0.3		
Diarrhea and gastroenteritis of infectious origin (A09)															
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	
Tuberculosis (A16-A19)															
2000.....	0.3	0.4	0.2	0.2	0.2	0.1	1.0	1.5	0.7	0.9	1.5	0.6			
1999.....	0.3	0.5	0.2	0.2	0.3	0.1	1.1	1.7	0.7	1.0	1.5	0.6			
Respiratory tuberculosis (A16)															
2000.....	0.2	0.3	0.1	0.1	0.2	0.1	0.8	1.2	0.5	0.8	1.3	0.4			
1999.....	0.3	0.4	0.2	0.2	0.3	0.1	0.9	1.4	0.5	0.8	1.2	0.5			
Other tuberculosis (A17-A19)															
2000.....	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	
1999.....	0.1	0.1	0.0	0.0	0.1	0.0	0.2	0.3	0.1	0.2	0.3	0.2	0.3	*	
Zoonotic and other bacterial diseases (A20-A49)															
2000.....	11.9	12.9	11.1	10.7	11.6	10.0	20.6	22.4	19.3	25.6	28.4	23.7			
1999.....	12.1	13.3	11.2	10.8	11.8	10.0	21.5	24.1	19.6	26.2	29.9	23.5			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Plague (A20)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Brucellosis (A23)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Tetanus (A33-A35)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Diphtheria (A36)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Whooping cough (A37)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Scarlet fever and erysipelas (A38,A46)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Meningococcal infection (A39)															
2000.....	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	*	*	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
Septicemia (A40-A41)															
2000.....	11.4	12.2	10.6	10.2	11.0	9.5	19.9	21.4	18.7	24.8	27.3	23.0			
1999.....	11.3	12.4	10.5	10.1	11.0	9.4	20.4	22.8	18.6	24.9	28.2	22.5			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION,
INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) - - ALL RACES - - - - - WHITE - - - - - TOTAL - - - - - ALL OTHER - - - - - BLACK - - -
AND YEAR BOTH SEXES MALE FEMALE SEXES MALE FEMALE SEXES MALE FEMALE SEXES MALE FEMALE

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Yellow fever (A95)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Other and unspecified arthropod-borne viral and hemorrhagic fevers (A90-A94,A96-A99)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Herpesviral (herpes simplex) infections (B00)															
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	0.1	*	*	*	*	
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	0.1	*	*	*	*	
Zoster (herpes zoster) (B02)															
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	
1999.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*	*	*	*	*	*	
Smallpox (B03)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Measles (B05)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Rubella (German measles) (B06)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Viral hepatitis (B15-B19)															
2000.....	1.9	2.6	1.3	1.8	2.4	1.2	2.9	3.9	2.1	2.7	3.9	1.8	1.8	1.8	
1999.....	1.8	2.4	1.2	1.6	2.3	1.1	2.7	3.5	2.0	2.6	3.5	1.9	1.9	1.9	

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER		
	BOTH SEXES		MALE									
	BOTH SEXES	MALE	FEMALE									
Human immunodeficiency virus (HIV) disease (B20-B24)												
2000.....	5.3	8.0	2.6	2.8	4.7	1.0	17.1	25.8	9.7	23.7	35.9	13.4
1999.....	5.4	8.4	2.6	3.0	5.0	1.0	17.6	26.9	9.7	24.2	37.1	13.4
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)												
2000.....	2.1	3.1	1.0	1.1	1.9	0.4	6.6	9.9	3.8	9.1	13.7	5.3
1999.....	2.2	3.3	1.1	1.2	2.0	0.5	7.0	10.6	3.9	9.5	14.5	5.3
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)												
2000.....	0.4	0.6	0.1	0.3	0.5	0.0	0.9	1.6	0.4	1.3	2.2	0.5
1999.....	0.4	0.7	0.1	0.3	0.5	0.1	1.0	1.7	0.4	1.3	2.3	0.5
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)												
2000.....	0.6	0.9	0.2	0.3	0.6	0.1	1.7	2.7	0.9	2.4	3.7	1.2
1999.....	0.6	1.0	0.2	0.4	0.6	0.1	1.7	2.7	0.8	2.3	3.7	1.2
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)												
2000.....	1.2	1.8	0.7	0.6	0.9	0.2	4.4	6.5	2.7	6.1	9.1	3.7
1999.....	1.3	1.9	0.7	0.6	0.9	0.2	4.7	7.1	2.8	6.5	9.8	3.8
Unspecified human immunodeficiency virus (HIV) disease (B24)												
2000.....	1.0	1.5	0.5	0.5	0.8	0.2	3.4	5.2	1.9	4.8	7.2	2.7
1999.....	1.0	1.5	0.5	0.5	0.9	0.2	3.3	4.9	1.8	4.5	6.8	2.5
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)												
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.4
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)												
2000.....	1.1	1.3	0.9	1.0	1.2	0.8	1.5	2.0	1.2	1.7	2.2	1.3
1999.....	1.5	1.8	1.3	1.3	1.6	1.1	2.5	3.0	2.2	2.9	3.3	2.6

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Mycoses (B35-B49)															
2000.....	0.4	0.6	0.3	0.4	0.5	0.3	0.7	1.0	0.6	0.8	1.1	0.7			
1999.....	0.7	0.9	0.5	0.6	0.8	0.4	1.2	1.4	1.0	1.3	1.6	1.2			
Protozoal diseases (B50-B64)															
2000.....	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.1	0.2	0.3	0.2			
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.3	0.2	0.3			
Malaria (B50-B54)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*			
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			
Leishmaniasis (B55)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*			
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			
Trypanosomiasis (B56-B57)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*			
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			
Pneumocystosis (B59)															
2000.....	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.1	0.2	0.2	0.1			
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.2	0.2			
Other and unspecified protozoal diseases (B58,B60-B64)															
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*			
1999.....	0.0	0.0	*	*	*	*	*	*	*	*	*	*			
Helminthiases (B65-B83)															
2000.....	0.0	0.0	*	0.0	*	*	*	*	*	*	*	*			
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE									
Schistosomiasis (bilharziasis) (B65)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Other and unspecified helminthiases (B66-B83)															
2000.....	0.0	0.0	*	0.0	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Sequelae of tuberculosis (B90)															
2000.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*	*	*	*	*	*	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	*	*	*	*	
Sequelae of poliomyelitis (B91)															
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	*	*	*	*	*	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	*	*	*	*	*	
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)															
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.6	0.4	0.5	0.7	0.4			
1999.....	0.5	0.6	0.5	0.4	0.5	0.4	1.0	1.1	0.9	1.2	1.4	1.1			
II. Neoplasms (C00-D48)															
2000.....	205.9	253.4	173.8	203.4	248.5	173.0	220.6	287.8	176.4	254.2	338.5	200.8			
1999.....	207.5	257.7	174.0	204.7	252.6	172.7	224.9	293.5	180.0	259.3	345.8	204.7			
Malignant neoplasms (C00-C97)															
2000.....	201.0	247.4	169.6	198.4	242.5	168.7	216.4	283.0	172.8	249.6	333.2	196.6			
1999.....	202.7	251.6	169.9	199.8	246.5	168.6	220.7	288.8	176.1	254.4	340.5	200.2			
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)															
2000.....	2.7	4.1	1.6	2.5	3.7	1.6	3.7	6.3	1.7	4.1	7.4	1.8			
1999.....	2.8	4.2	1.6	2.6	3.8	1.6	3.8	6.4	1.9	4.4	7.5	2.1			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Of lip (C00)															
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*	*	*	*	*	*	*
Of tongue (C01-C02)															
2000.....	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.1	0.3	0.8	1.4	0.3			
1999.....	0.6	1.0	0.4	0.6	0.9	0.4	0.7	1.1	0.4	0.9	1.4	0.4			
Of pharynx (C10-C13,C14.0)															
2000.....	1.1	1.8	0.6	1.0	1.5	0.5	2.1	3.7	0.9	2.2	4.1	0.9			
1999.....	1.1	1.8	0.6	1.0	1.6	0.5	2.1	3.6	0.9	2.3	4.1	1.0			
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)															
2000.....	0.9	1.3	0.6	0.9	1.3	0.6	1.0	1.6	0.5	1.1	1.9	0.5			
1999.....	1.0	1.3	0.6	0.9	1.3	0.7	1.0	1.7	0.6	1.2	2.0	0.7			
Malignant neoplasms of digestive organs (C15-C26)															
2000.....	47.5	60.2	37.6	45.6	57.8	36.0	60.5	76.8	48.6	66.5	84.8	53.5			
1999.....	47.7	60.7	37.8	45.8	58.4	36.2	60.8	77.1	48.8	67.1	86.1	53.7			
Of esophagus (C15)															
2000.....	4.4	7.7	1.8	4.3	7.6	1.7	5.3	8.7	2.7	6.5	10.9	3.3			
1999.....	4.4	7.7	1.8	4.2	7.5	1.7	5.4	9.4	2.5	6.6	11.5	3.1			
Of stomach (C16)															
2000.....	4.6	6.4	3.3	4.0	5.6	2.8	8.9	12.2	6.7	9.0	12.5	6.7			
1999.....	4.7	6.6	3.3	4.1	5.8	2.8	9.1	12.7	6.6	9.2	13.3	6.6			
Of small intestine (C17)															
2000.....	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.6	0.4	0.5	0.6	0.4			
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.5	0.7	0.4	0.6	0.8	0.5			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES		MALE	BOTH SEXES		MALE									
Of colon, rectum and anus (C18-C21)															
2000.....	20.9	24.9	17.9	20.4	24.3	17.4	24.3	29.2	20.9	28.3	34.1	24.4			
1999.....	21.1	25.2	18.1	20.6	24.8	17.5	24.4	28.3	21.6	28.6	33.6	25.1			
Colon (C18)															
2000.....	17.7	20.8	15.3	17.1	20.2	14.8	21.0	24.8	18.3	24.7	29.2	21.7			
1999.....	17.9	21.2	15.5	17.4	20.8	14.9	21.1	23.9	19.1	25.0	28.7	22.4			
Rectosigmoid junction and rectum (C19-C20)															
2000.....	3.1	4.0	2.3	3.0	3.9	2.3	3.2	4.2	2.4	3.5	4.6	2.6			
1999.....	3.1	3.9	2.4	3.0	3.8	2.4	3.1	4.2	2.4	3.4	4.7	2.5			
Anus and anal canal (C21)															
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2			
1999.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2			
Of liver and intrahepatic bile ducts (C22)															
2000.....	4.7	6.8	3.0	4.3	6.2	2.7	7.4	11.2	4.5	6.1	9.5	3.6			
1999.....	4.6	6.5	3.0	4.2	6.0	2.8	7.0	10.2	4.5	6.0	8.7	3.9			
Liver (C22.0,C22.2-C22.9)															
2000.....	3.7	5.7	2.1	3.3	5.0	1.9	6.4	10.1	3.6	5.3	8.6	2.9			
1999.....	3.6	5.5	2.1	3.3	4.9	1.9	6.1	9.1	3.6	5.2	7.9	3.2			
Intrahepatic bile duct carcinoma (C22.1)															
2000.....	1.0	1.2	0.9	1.0	1.2	0.9	1.0	1.1	0.9	0.8	0.9	0.8			
1999.....	0.9	1.1	0.9	0.9	1.1	0.9	0.9	1.0	0.8	0.8	0.8	0.7			
Of gallbladder and extrahepatic bile ducts (C23-C24)															
2000.....	1.3	1.2	1.4	1.3	1.2	1.4	1.3	1.1	1.4	1.2	0.9	1.4			
1999.....	1.3	1.2	1.4	1.3	1.2	1.4	1.4	1.1	1.6	1.3	1.0	1.5			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Malignant neoplasms of bone and articular cartilage (C40-C41)															
2000.....	0.5	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.3	0.4	0.6	0.3			
1999.....	0.4	0.5	0.4	0.5	0.5	0.4	0.4	0.5	0.3	0.4	0.5	0.3			
Melanoma and other malignant neoplasms of skin (C43-C44)															
2000.....	3.5	5.2	2.3	3.9	5.7	2.5	1.0	1.3	0.8	1.1	1.5	0.9			
1999.....	3.5	5.3	2.2	3.9	5.8	2.5	0.9	1.4	0.5	1.0	1.6	0.6			
Malignant melanoma of skin (C43)															
2000.....	2.7	3.8	1.8	3.0	4.3	2.1	0.5	0.5	0.5	0.6	0.6	0.5			
1999.....	2.7	3.8	1.8	3.0	4.3	2.0	0.4	0.5	0.3	0.4	0.5	0.3			
Other malignant neoplasms of skin (C44)															
2000.....	0.8	1.4	0.4	0.8	1.4	0.4	0.5	0.8	0.3	0.6	0.9	0.4			
1999.....	0.8	1.4	0.4	0.9	1.5	0.5	0.5	0.8	0.2	0.6	1.1	0.2			
Malignant neoplasms of mesothelial and soft tissue (C45-C49)															
2000.....	2.5	3.3	1.8	2.6	3.5	1.9	1.9	2.2	1.6	2.1	2.5	1.9			
1999.....	2.5	3.3	1.8	2.5	3.5	1.8	2.0	2.3	1.7	2.2	2.6	1.9			
Mesothelioma (C45)															
2000.....	0.9	1.7	0.3	0.9	1.8	0.3	0.4	0.7	0.1	0.4	0.8	0.1			
1999.....	0.9	1.6	0.3	0.9	1.8	0.3	0.4	0.7	0.2	0.4	0.8	0.2			
Kaposi's sarcoma (C46)															
2000.....	0.0	0.0	*	0.0	0.0	*	0.0	0.1	*	*	*	*			
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*	*	*	*			
Of other mesothelial and soft tissue (C47-C49)															
2000.....	1.6	1.6	1.6	1.6	1.7	1.6	1.5	1.4	1.5	1.6	1.6	1.7			
1999.....	1.6	1.7	1.5	1.6	1.7	1.5	1.6	1.6	1.6	1.8	1.8	1.7			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Malignant neoplasm of breast (C50)															
2000.....	15.4	0.4	27.1	15.0	0.4	26.6	17.1	0.5	29.2	20.7	0.6	34.9			
1999.....	15.3	0.3	27.0	14.9	0.3	26.4	17.5	0.5	29.9	21.1	0.7	35.6			
Malignant neoplasms of female genital organs (C51-C58)															
2000.....	9.6	...	17.0	9.4	...	16.8	10.8	...	18.4	12.5	...	21.1			
1999.....	9.5	...	16.9	9.4	...	16.7	10.6	...	18.1	12.4	...	20.9			
Of vulva and vagina (C51-C52)															
2000.....	0.4	...	0.7	0.4	...	0.7	0.3	...	0.6	0.4	...	0.7			
1999.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.6	0.4	...	0.7			
Of cervix uteri (C53)															
2000.....	1.5	...	2.8	1.3	...	2.5	2.8	...	4.8	3.2	...	5.5			
1999.....	1.6	...	2.9	1.4	...	2.6	2.7	...	4.7	3.2	...	5.5			
Of corpus uteri and uterus, part unspecified (C54-C55)															
2000.....	2.4	...	4.2	2.2	...	3.9	3.6	...	6.1	4.4	...	7.3			
1999.....	2.4	...	4.1	2.3	...	3.9	3.4	...	5.7	4.2	...	6.9			
Of ovary (C56)															
2000.....	5.1	...	9.0	5.3	...	9.4	3.9	...	6.7	4.3	...	7.3			
1999.....	5.0	...	8.9	5.2	...	9.2	4.0	...	6.8	4.4	...	7.5			
Of other and unspecified female genital organs (C57)															
2000.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.3	0.2	...	0.3			
1999.....	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3	0.2	...	0.3			
Of placenta (C58)															
2000.....	*	...	*	*	...	*	*	...	*	*	...	*		*	
1999.....	*	...	*	*	...	*	*	...	*	*	...	*		*	

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Malignant neoplasms of male genital organs (C60-C63)															
2000.....	11.5	30.2	...	10.5	27.7	...	19.4	51.2	...	24.2	65.7	...			
1999.....	11.9	31.4	...	11.0	28.9	...	19.8	52.5	...	24.5	66.9	...			
Of prostate (C61)															
2000.....	11.3	29.7	...	10.3	27.2	...	19.3	50.9	...	24.0	65.3	...			
1999.....	11.7	30.9	...	10.7	28.4	...	19.6	52.1	...	24.3	66.5	...			
Of testis (C62)															
2000.....	0.1	0.3	...	0.1	0.3	...	0.1	0.1	...	0.1	0.2	...			
1999.....	0.2	0.3	...	0.2	0.3	...	0.1	0.1	...	0.1	0.1	...			
Of penis and other and unspecified male genital organs (C60,C63)															
2000.....	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...			
1999.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...	0.1	0.3	...			
Malignant neoplasms of urinary tract (C64-C68)															
2000.....	8.8	13.9	5.3	9.1	14.4	5.4	7.0	10.1	4.9	8.1	11.8	5.7			
1999.....	8.7	13.8	5.2	8.9	14.1	5.2	7.3	10.3	5.3	8.4	11.8	6.1			
Of kidney and renal pelvis (C64-C65)															
2000.....	4.3	6.1	2.9	4.3	6.2	2.9	3.7	5.4	2.5	4.2	6.2	2.8			
1999.....	4.1	5.9	2.7	4.1	6.0	2.7	3.7	5.3	2.5	4.2	6.0	2.9			
Of bladder (C67)															
2000.....	4.3	7.5	2.3	4.5	7.8	2.3	3.1	4.6	2.3	3.8	5.4	2.8			
1999.....	4.4	7.6	2.3	4.5	7.9	2.3	3.4	4.8	2.5	4.0	5.6	3.0			
Of other and unspecified sites within the urinary tract (C66,C68)															
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.1	0.2	0.1	0.2	*	0.2			
1999.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	*	0.2			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	*	*	*	*	*	*	*
Malignant neoplasm of eye and adnexa (C69)																
2000	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*	*	*	*	*	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	*	*	*	*	*	*	*
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)																
2000	4.6	5.6	3.7	4.9	6.0	4.0	2.5	3.0	2.2	2.7	3.3	2.3				
1999	4.7	5.7	3.9	5.0	6.1	4.2	2.5	3.0	2.1	2.8	3.3	2.4				
Of brain (C71)																
2000	4.5	5.5	3.6	4.8	5.9	3.9	2.5	2.9	2.1	2.6	3.2	2.2				
1999	4.6	5.6	3.8	4.9	6.0	4.1	2.4	2.9	2.0	2.7	3.2	2.2				
Of other parts of central nervous system (C70,C72)																
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	*	*	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	*
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)																
2000	0.8	0.8	0.7	0.7	0.8	0.7	0.8	0.6	0.9	0.8	0.6	0.9				
1999	0.7	0.8	0.7	0.7	0.8	0.7	0.7	0.7	0.8	0.8	0.7	0.7				
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)																
2000	14.8	17.7	12.7	14.5	17.2	12.6	16.3	20.7	13.2	18.8	24.4	15.0				
1999	16.2	19.5	13.7	15.9	19.1	13.5	18.0	22.2	15.0	20.8	26.1	17.1				
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)																
2000	20.5	25.8	16.7	20.8	26.3	16.8	17.8	21.6	15.1	20.1	24.6	16.9				
1999	20.7	26.1	16.9	21.0	26.5	17.1	18.2	22.5	15.3	20.4	25.3	17.1				
Hodgkin's disease (C81)																
2000	0.5	0.6	0.4	0.5	0.6	0.4	0.3	0.4	0.3	0.3	0.4	0.3				
1999	0.5	0.7	0.4	0.5	0.7	0.4	0.3	0.4	0.3	0.3	0.4	0.3				

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)															
2000.....	8.3	10.2	6.8	8.7	10.7	7.2	5.3	6.5	4.3	5.3	6.6	4.3			
1999.....	8.4	10.4	7.0	8.8	10.7	7.4	5.5	7.2	4.3	5.7	7.4	4.5			
Non-Hodgkin's lymphoma (C82-C85)															
2000.....	8.3	10.2	6.8	8.6	10.6	7.1	5.2	6.5	4.3	5.3	6.5	4.3			
1999.....	8.4	10.3	7.0	8.8	10.7	7.3	5.5	7.2	4.3	5.7	7.4	4.5			
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)															
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*			
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*			
Malignant immunoproliferative diseases (C88)															
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	*	*	*	*			
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	*	*	0.1	*	*			
Multiple myeloma and malignant plasma cell neoplasms (C90)															
2000.....	3.9	4.7	3.4	3.6	4.4	3.0	6.1	6.7	5.6	7.6	8.6	6.9			
1999.....	3.9	4.7	3.3	3.6	4.4	3.0	6.2	7.0	5.7	7.5	8.6	6.8			
Leukemia (C91-C95)															
2000.....	7.8	10.2	6.0	7.9	10.5	6.1	6.1	7.8	4.8	6.7	8.8	5.3			
1999.....	7.7	10.2	6.0	7.9	10.4	6.2	6.1	7.8	5.0	6.7	8.5	5.5			
Lymphoid leukemia (C91)															
2000.....	2.2	3.1	1.6	2.3	3.2	1.7	1.6	2.3	1.1	1.9	2.8	1.3			
1999.....	2.3	3.2	1.7	2.4	3.3	1.7	1.7	2.3	1.3	2.0	2.9	1.5			
Myeloid leukemia (C92)															
2000.....	3.5	4.6	2.8	3.6	4.7	2.9	2.8	3.3	2.4	3.1	3.6	2.7			
1999.....	3.4	4.3	2.8	3.5	4.4	2.8	2.8	3.4	2.3	2.9	3.6	2.4			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Monocytic leukemia (C93)															
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*	*	*	*	*	*	*
Other and unspecified leukemia (C94-C95)															
2000.....	1.9	2.5	1.5	1.9	2.5	1.6	1.6	2.1	1.2	1.7	2.4	1.3			
1999.....	2.0	2.5	1.5	2.0	2.6	1.6	1.6	2.0	1.4	1.8	2.1	1.6			
In situ neoplasms (D00-D09)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Carcinoma in situ of breast and genitourinary system (D05-D07)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Carcinoma in situ of other and unspecified sites (D00-D04,D09)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
Benign neoplasms (D10-D36)															
2000.....	0.6	0.5	0.6	0.5	0.5	0.6	0.7	0.7	0.8	0.9	0.7	1.0			
1999.....	0.6	0.6	0.6	0.5	0.5	0.5	0.8	0.8	0.8	1.0	0.9	1.0			
Of female genital organs (D25-D28)															
2000.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1	0.1	...	0.2			
1999.....	0.0	...	0.0	*	...	*	*	...	*	*	...	*			
Of eye, brain and other parts of central nervous system (D31-D33)															
2000.....	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5	0.5	0.4	0.5			
1999.....	0.4	0.3	0.4	0.3	0.3	0.3	0.5	0.4	0.5	0.6	0.5	0.6			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Of other and unspecified sites (D10-D24, D29-D30, D34-D36)															
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2	0.3	0.4	0.4	0.3	0.3	0.3
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)															
2000.....	4.1	5.2	3.4	4.2	5.4	3.5	3.2	4.0	2.6	3.5	4.2	3.0			
1999.....	4.1	5.3	3.3	4.2	5.5	3.4	3.2	3.7	2.8	3.6	4.1	3.2			
Neoplasm of uncertain or unknown behavior of unspecified sites (D48)															
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.3	0.2	0.2
1999.....	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.2	0.3	0.2	0.3	0.2	0.3
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)															
2000.....	3.4	3.4	3.4	2.9	3.0	2.8	5.9	5.8	6.0	7.4	7.2	7.7			
1999.....	3.3	3.5	3.3	2.9	3.0	2.8	6.0	6.0	6.0	7.4	7.4	7.4			
Anemias (D50-D64)															
2000.....	1.6	1.5	1.7	1.4	1.4	1.4	2.8	2.7	2.8	3.5	3.4	3.5			
1999.....	1.6	1.7	1.6	1.4	1.4	1.4	3.0	3.0	2.9	3.7	3.7	3.6			
Nutritional anemias (D50-D53)															
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.2	*	0.2			
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.2	*	0.2			
Hemolytic anemias (D55-D59)															
2000.....	0.3	0.3	0.3	0.1	0.1	0.1	1.0	1.0	1.1	1.5	1.3	1.5			
1999.....	0.3	0.3	0.3	0.1	0.1	0.1	1.1	1.3	1.0	1.6	1.8	1.4			
Aplastic anemias (D60-D61)															
2000.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.4	0.5	0.4			
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.5	0.5	0.4			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Acute posthemorrhagic and other anemias (D62-D64)															
2000.....	0.8	0.8	0.8	0.8	0.7	0.8	1.2	1.1	1.2	1.4	1.3	1.5			
1999.....	0.8	0.8	0.9	0.8	0.7	0.8	1.3	1.1	1.3	1.5	1.3	1.5			
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)															
2000.....	1.0	1.0	0.9	0.9	1.0	0.9	1.3	1.3	1.3	1.5	1.5	1.6			
1999.....	1.0	1.0	0.9	0.9	0.9	0.8	1.4	1.5	1.3	1.6	1.7	1.6			
Other diseases of blood and blood-forming organs (D70-D76)															
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3	0.4	0.3			
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3			
Certain disorders involving the immune mechanism (D80-D89)															
2000.....	0.5	0.5	0.5	0.3	0.3	0.3	1.5	1.4	1.7	2.1	1.9	2.3			
1999.....	0.4	0.4	0.4	0.3	0.3	0.2	1.3	1.1	1.4	1.8	1.5	1.9			
IV. Endocrine, nutritional and metabolic diseases (E00-E88)															
2000.....	34.3	36.6	32.2	31.7	34.6	29.2	52.6	50.6	53.2	62.4	60.1	63.0			
1999.....	34.2	36.5	32.2	31.6	34.3	29.1	53.8	53.0	53.8	62.7	61.9	62.6			
Endocrine diseases (E00-E34)															
2000.....	26.1	28.2	24.4	23.8	26.4	21.7	43.3	41.5	43.9	51.2	48.9	52.0			
1999.....	26.2	28.4	24.5	23.7	26.4	21.6	44.5	43.0	45.0	51.6	50.0	52.0			
Disorders of thyroid gland (E00-E07)															
2000.....	0.6	0.4	0.8	0.6	0.4	0.8	0.7	0.4	0.8	0.8	0.6	1.0			
1999.....	0.7	0.4	0.8	0.6	0.4	0.8	0.7	0.5	0.8	0.9	0.6	1.0			
Diabetes mellitus (E10-E14)															
2000.....	25.2	27.6	23.3	22.8	25.8	20.6	42.0	40.6	42.5	49.7	47.8	50.4			
1999.....	25.2	27.7	23.3	22.8	25.8	20.5	43.2	41.8	43.7	50.1	48.6	50.4			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Disorders of pituitary gland (E22-E23)															
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	*	0.2	0.2	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1
Disorders of adrenal glands (E24-E27)															
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	*	*	0.2	*	0.2
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	*	0.1	*	*
Diseases of thymus (E32)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other endocrine diseases (E15-E21,E28-E31,E34)															
2000.....	0.2	0.1	0.2	0.1	0.1	0.1	0.3	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.4
1999.....	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.4	0.2	0.4	0.4	0.5	0.5	0.3	0.3
Nutritional diseases (E40-E68)															
2000.....	2.6	2.5	2.7	2.5	2.3	2.5	3.6	3.2	3.8	4.4	4.1	4.5	4.4	4.1	4.5
1999.....	2.5	2.4	2.6	2.4	2.3	2.4	3.6	3.6	3.6	4.4	4.4	4.4	4.4	4.4	4.4
Malnutrition (E40-E46)															
2000.....	1.4	1.3	1.5	1.4	1.2	1.4	1.9	1.9	1.9	2.3	2.4	2.1	2.4	2.1	2.1
1999.....	1.5	1.4	1.5	1.4	1.2	1.4	2.1	2.4	1.9	2.5	2.5	2.8	2.3	2.3	2.3
Nutritional marasmus (E41)															
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	*	0.1	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	*	0.1	*	*
Other and unspecified malnutrition (E40,E42-E46)															
2000.....	1.3	1.2	1.4	1.3	1.1	1.3	1.8	1.8	1.8	2.2	2.2	2.2	2.1	2.1	2.1
1999.....	1.4	1.3	1.4	1.3	1.2	1.4	2.0	2.3	1.8	2.3	2.3	2.7	2.7	2.1	2.1

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	*	0.1	0.1	
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2	0.2	0.2	*	0.2	0.2		
Other nutritional deficiencies (E50-E64)																
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	*	0.1	0.1		
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2	0.2	0.2	*	0.2	0.2		
Obesity and other hyperalimentation (E65-E68)																
2000	1.1	1.1	1.1	1.0	1.0	1.0	1.5	1.2	1.7	2.0	1.6	2.2				
1999	1.0	0.9	1.0	0.9	0.9	0.8	1.3	1.1	1.5	1.7	1.4	2.0				
Metabolic disorders (E70-E88)																
2000	5.5	5.9	5.0	5.5	5.9	5.0	5.8	5.8	5.5	6.8	7.1	6.5				
1999	5.5	5.7	5.2	5.5	5.6	5.2	5.7	6.4	5.2	6.7	7.5	6.1				
Cystic fibrosis (E84)																
2000	0.2	0.2	0.2	0.2	0.2	0.2	0.0	*	*	*	*	*	*	*		
1999	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*	*	*	*	*	*		
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)																
2000	2.8	2.6	2.9	2.7	2.5	2.8	3.5	3.4	3.5	4.3	4.3	4.1				
1999	3.0	2.7	3.1	2.9	2.5	3.1	3.6	4.1	3.2	4.3	5.0	3.8				
Other metabolic disorders (E70-E83,E85,E88)																
2000	2.5	3.0	2.0	2.5	3.1	2.0	2.2	2.4	2.0	2.5	2.7	2.3				
1999	2.3	2.8	1.9	2.3	2.8	1.9	2.1	2.3	2.0	2.4	2.5	2.3				
V. Mental and behavioral disorders (F01-F99)																
2000	16.7	16.4	15.9	16.7	16.2	16.1	15.3	17.8	12.9	18.0	21.6	14.9				
1999	15.6	15.9	14.5	15.7	15.6	14.7	13.8	16.9	11.0	16.1	20.1	12.7				
Organic dementia (F01,F03)																
2000	11.9	9.9	12.7	12.1	10.0	13.0	9.7	9.0	9.9	11.1	10.7	11.2				
1999	10.6	9.1	11.2	10.9	9.3	11.6	7.7	7.3	7.7	8.7	8.4	8.7				

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Other and unspecified organic mental disorders (F04-F09)															
2000.....	0.8	0.7	0.9	0.8	0.7	0.9	0.6	0.7	0.6	0.8	0.8	0.8	0.7		
1999.....	0.9	0.8	0.9	0.9	0.8	1.0	0.7	0.8	0.7	0.9	1.0	0.8			
Mental and behavioral disorders due to psychoactive substance use (F10-F19)															
2000.....	3.2	5.1	1.4	2.9	4.7	1.4	4.4	7.6	1.9	5.5	9.5	2.4			
1999.....	3.2	5.1	1.4	2.9	4.7	1.2	4.8	8.1	2.1	5.8	9.9	2.6			
Mental and behavioral disorders due to use of alcohol (F10)															
2000.....	2.3	3.8	0.9	2.1	3.5	0.8	3.1	5.6	1.1	3.6	6.7	1.3			
1999.....	2.3	3.8	0.9	2.1	3.5	0.8	3.3	5.8	1.3	3.9	6.9	1.5			
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)															
2000.....	0.9	1.2	0.6	0.8	1.1	0.5	1.3	2.0	0.8	1.8	2.7	1.1			
1999.....	0.9	1.3	0.5	0.8	1.2	0.4	1.5	2.3	0.9	2.0	3.0	1.1			
Schizophrenia, schizotypal and delusional disorders (F20-F29)															
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.2		
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2		
Mood (affective) disorders (F30-F39)															
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	0.1	0.1	*	0.1	*	0.1	*
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	*	0.1	*	0.1	*	*	*
Neurotic, stress-related and somatoform disorders (F40-F48)															
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*
Mental retardation (F70-F79)															
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1		
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	0.2	0.2		

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Other and unspecified mental and behavioral disorders (F50-F69, F80-F99)															
2000.....	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	0.1	0.1	*	*	*	*	
1999.....	0.1	0.1	0.2	0.2	0.1	0.2	0.1	*	*	0.1	*	0.1	*	*	
VI. Diseases of the nervous system (G00-G98)															
2000.....	33.0	34.1	31.8	34.0	35.1	32.7	23.5	24.2	22.8	27.3	27.9	26.4			
1999.....	31.1	32.8	29.5	32.0	33.8	30.3	22.0	22.7	21.3	25.2	26.0	24.4			
Meningitis (G00,G03)															
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.5	0.6	0.8	0.6			
1999.....	0.3	0.3	0.3	0.3	0.3	0.2	0.6	0.7	0.5	0.7	0.9	0.6			
Other inflammatory diseases of central nervous system (G04-G09)															
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.3			
1999.....	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.4	0.3			
Parkinson's disease (G20-G21)															
2000.....	5.7	8.6	4.0	6.0	9.1	4.2	2.6	4.2	1.7	2.5	4.0	1.6			
1999.....	5.4	8.2	3.7	5.7	8.7	3.9	2.5	3.8	1.9	2.4	3.6	1.7			
Alzheimer's disease (G30)															
2000.....	18.0	14.7	19.5	18.7	15.4	20.3	11.2	9.2	12.3	13.1	10.8	14.2			
1999.....	16.5	14.0	17.6	17.2	14.7	18.4	9.9	8.3	10.7	11.4	9.7	12.2			
Multiple sclerosis (G35)															
2000.....	1.0	0.8	1.2	1.1	0.8	1.3	0.8	0.5	1.0	1.0	0.7	1.3			
1999.....	1.0	0.7	1.2	1.0	0.7	1.3	0.6	0.4	0.8	0.8	0.6	1.1			
Epilepsy (G40-G41)															
2000.....	0.5	0.5	0.4	0.4	0.5	0.4	0.8	1.0	0.7	1.0	1.3	0.9			
1999.....	0.5	0.6	0.4	0.4	0.5	0.4	0.9	1.0	0.7	1.1	1.4	0.9			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	1.3	0.9	1.5	1.3	0.9	1.6	1.0	0.7	1.2	1.0	0.7	1.2	1.0	0.7	1.2
Chronic rheumatic heart diseases (I05-I09)	1999	1.3	0.9	1.6	1.4	0.9	1.7	1.1	0.7	1.3	1.0	0.7	1.2	1.0	0.7	1.2
Rheumatic mitral valve diseases (I05)	2000	0.7	0.5	0.9	0.8	0.5	0.9	0.5	0.4	0.6	0.5	0.4	0.6	0.5	0.4	0.6
	1999	0.7	0.5	1.0	0.8	0.5	0.9	0.6	0.4	0.8	0.6	0.5	0.6	0.5	0.5	0.8
Rheumatic aortic valve diseases (I06)	2000	0.0	*	*	0.0	*	*	*	*	*	*	*	*	*	*	*
	1999	0.0	*	0.0	0.0	*	0.0	*	*	*	*	*	*	*	*	*
Disorders of both mitral and aortic valves (I08.0)	2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	*	*	*	*
	1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	*	*	*	*	*
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)	2000	0.4	0.3	0.5	0.4	0.3	0.5	0.4	0.2	0.5	0.4	0.2	0.5	0.4	0.2	0.5
	1999	0.4	0.3	0.5	0.5	0.3	0.6	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.4
Hypertensive diseases (I10-I13)	2000	16.2	16.1	15.6	13.5	13.2	13.1	35.3	36.8	33.4	43.6	46.3	40.8			
	1999	15.9	15.8	15.4	13.2	12.8	12.8	35.5	37.5	33.4	43.5	46.8	40.3			
Essential (primary) hypertension (I10)	2000	3.7	3.3	3.8	3.4	3.0	3.5	6.2	6.1	6.0	7.4	7.4	7.1			
	1999	3.7	3.2	3.8	3.3	2.9	3.5	6.2	6.1	6.1	7.3	7.2	7.2			
Hypertensive heart disease (I11)	2000	8.6	8.6	8.1	7.1	7.0	6.8	19.0	20.3	17.5	23.8	25.9	21.6			
	1999	8.4	8.3	8.0	6.9	6.7	6.7	18.6	20.1	17.1	23.1	25.5	21.0			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES				WHITE				TOTAL				ALL OTHER			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
Hypertensive renal disease (I12)																
2000.....	2.8	3.0	2.7	2.2	2.4	2.0	7.4	7.4	7.3	9.1	9.3	8.8				
1999.....	2.6	2.9	2.4	2.0	2.2	1.8	7.3	7.7	6.9	8.8	9.6	8.2				
Hypertensive heart and renal disease (I13)																
2000.....	1.0	1.1	1.0	0.8	0.8	0.7	2.8	2.9	2.6	3.4	3.7	3.2				
1999.....	1.2	1.3	1.1	0.9	1.0	0.9	3.5	3.6	3.3	4.2	4.4	4.0				
Ischemic heart diseases (I20-I25)																
2000.....	186.9	237.7	147.9	185.7	238.0	145.3	190.7	229.6	161.7	219.3	262.4	187.5				
1999.....	195.6	249.0	154.9	194.5	249.4	152.5	198.1	239.5	167.4	226.4	272.6	192.5				
Acute myocardial infarction (I21-I22)																
2000.....	70.0	90.1	54.7	69.8	90.7	53.6	70.8	84.5	60.8	81.4	96.3	70.7				
1999.....	73.6	95.4	57.1	73.5	96.3	56.2	72.9	87.3	62.5	83.7	99.8	72.2				
Other acute ischemic heart diseases (I24)																
2000.....	1.2	1.5	0.9	1.2	1.4	0.9	1.6	2.0	1.2	1.9	2.5	1.4				
1999.....	1.2	1.5	0.9	1.1	1.4	0.9	1.5	2.0	1.1	1.8	2.4	1.3				
Other forms of chronic ischemic heart disease (I20,I25)																
2000.....	115.7	146.1	92.3	114.8	145.9	90.8	118.4	143.0	99.7	136.1	163.7	115.4				
1999.....	120.8	152.1	96.9	119.8	151.7	95.4	123.7	150.3	103.8	140.9	170.4	119.0				
Atherosclerotic cardiovascular disease, so described (I25.0)																
2000.....	25.2	31.9	19.6	23.6	29.8	18.3	36.6	47.5	28.4	43.7	57.4	33.7				
1999.....	26.5	33.3	20.9	24.8	31.0	19.5	39.0	50.5	30.3	46.2	60.5	35.7				
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)																
2000.....	90.5	114.2	72.8	91.2	116.1	72.5	81.8	95.5	71.3	92.4	106.2	81.7				
1999.....	94.2	118.7	75.9	95.0	120.6	75.8	84.7	99.8	73.5	94.8	109.8	83.4				

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER		
	BOTH SEXES		MALE									
	BOTH SEXES	MALE	FEMALE									
Other heart diseases (I26-I51)												
2000.....	60.0	66.7	54.4	58.6	65.1	53.1	65.9	74.4	59.3	79.0	90.0	70.6
1999.....	61.2	68.5	55.2	59.7	66.6	53.8	68.8	79.4	60.9	81.9	95.7	71.8
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
2000.....	4.7	4.5	5.0	4.5	4.3	4.6	6.3	5.8	6.8	8.0	7.5	8.5
1999.....	4.9	4.7	5.0	4.6	4.5	4.6	6.9	6.5	7.3	8.8	8.3	9.1
Pulmonary embolism (I26)												
2000.....	3.1	3.0	3.1	2.9	2.9	2.9	4.3	4.2	4.3	5.5	5.6	5.5
1999.....	3.3	3.3	3.3	3.1	3.1	3.1	4.8	4.8	4.8	6.2	6.4	6.1
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
2000.....	1.7	1.4	1.9	1.6	1.4	1.7	2.1	1.5	2.5	2.5	1.9	3.0
1999.....	1.6	1.4	1.7	1.5	1.4	1.5	2.2	1.7	2.5	2.6	1.9	3.1
Other non-pulmonary forms of heart disease (I30-I51)												
2000.....	55.3	62.2	49.4	54.2	60.9	48.5	59.6	68.7	52.5	71.0	82.5	62.1
1999.....	56.3	63.8	50.2	55.1	62.1	49.2	61.9	72.9	53.6	73.1	87.4	62.7
Acute and subacute endocarditis (I33)												
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.5	0.8	0.9	0.7
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.7	0.5	0.7	0.9	0.6
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
2000.....	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.5	0.3	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.4	0.5	0.5	0.5
Other diseases of endocardium (I34-I38)												
2000.....	7.2	7.3	7.0	7.4	7.6	7.2	4.7	4.7	4.6	5.3	5.3	5.2
1999.....	7.2	7.5	7.0	7.5	7.7	7.2	4.9	5.0	4.8	5.3	5.3	5.3

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	1.0	0.9	1.1	1.1	1.0	1.2	0.7	0.5	0.8	0.8	0.6	0.9	1.1	0.9	0.9
Nonrheumatic mitral valve disorders (I34)																
2000.....	1.0	0.9	1.1	1.1	1.0	1.2	0.7	0.5	0.8	0.8	0.6	0.9	1.1	0.9	0.9	
1999.....	1.1	0.9	1.1	1.1	1.0	1.2	0.8	0.6	0.9	0.8	0.5	0.9	1.1	0.9	0.9	
Nonrheumatic aortic valve disorders (I35)																
2000.....	4.5	4.8	4.2	4.7	5.0	4.4	2.4	2.6	2.3	2.7	2.8	2.6	2.7	2.8	2.6	
1999.....	4.5	4.9	4.2	4.7	5.1	4.4	2.6	2.9	2.4	2.8	3.0	2.5	2.8	3.0	2.5	
All other diseases of endocardium (I36-I38)																
2000.....	1.6	1.6	1.6	1.6	1.5	1.6	1.6	1.6	1.5	1.8	1.9	1.8	1.9	1.8	1.8	
1999.....	1.7	1.6	1.7	1.7	1.6	1.7	1.6	1.6	1.5	1.8	1.8	1.8	1.8	1.8	1.8	
Cardiomyopathy (I42)																
2000.....	9.9	13.7	7.0	9.1	12.6	6.4	14.7	20.1	10.7	17.9	24.6	13.0	13.0	24.6	13.0	
1999.....	10.1	13.8	7.2	9.2	12.7	6.5	15.3	21.1	11.0	18.4	25.7	13.1	13.1	25.7	13.1	
Conduction disorders and cardiac dysrhythmias (I44-I49)																
2000.....	13.9	15.3	12.6	13.5	14.9	12.2	16.0	17.9	14.5	18.9	21.2	17.0	17.0	21.2	17.0	
1999.....	14.8	16.6	13.2	14.4	16.1	12.9	16.9	20.0	14.8	20.2	24.0	17.5	17.5	24.0	17.5	
Heart failure (I50)																
2000.....	20.2	20.9	19.4	20.3	21.0	19.4	18.6	19.2	18.1	22.1	23.2	21.3	21.3	23.2	21.3	
1999.....	20.3	21.0	19.5	20.3	21.0	19.5	19.3	20.3	18.5	22.7	24.4	21.4	21.4	24.4	21.4	
Congestive heart failure (I50.0)																
2000.....	18.7	19.3	18.0	18.8	19.4	18.1	16.7	17.0	16.4	19.8	20.4	19.3	19.3	20.4	19.3	
1999.....	18.8	19.4	18.2	18.9	19.4	18.2	17.4	18.3	16.7	20.5	21.9	19.4	19.4	21.9	19.4	
Other and unspecified heart failure (I50.1-I50.9)																
2000.....	1.5	1.7	1.4	1.4	1.6	1.3	1.9	2.2	1.7	2.3	2.7	2.0	2.0	2.7	2.0	
1999.....	1.5	1.7	1.4	1.4	1.6	1.3	1.9	2.1	1.7	2.2	2.5	2.0	2.0	2.5	2.0	

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.4	
1999	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3	0.4	0.4	0.5	0.4	0.3		
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)																
2000	3.2	3.8	2.6	3.0	3.7	2.5	4.1	5.1	3.4	5.1	6.4	4.1				
1999	3.0	3.7	2.4	2.8	3.5	2.3	3.9	4.7	3.3	4.7	5.9	3.8				
All other and ill-defined forms of heart disease (I51.0-I51.3, I51.6-I51.9)																
2000	60.8	61.1	59.7	58.7	58.6	57.8	73.9	77.9	70.3	82.4	87.1	78.1				
1999	61.8	62.4	60.5	59.8	60.0	58.7	74.1	78.4	70.4	82.4	87.4	78.1				
Cerebrovascular diseases (I60-I69)																
2000	2.4	1.8	2.9	2.2	1.7	2.6	3.3	2.2	4.1	3.4	2.3	4.3				
1999	2.4	1.8	2.8	2.3	1.7	2.6	3.2	2.2	4.0	3.3	2.4	4.0				
Subarachnoid hemorrhage (I60)																
2000	9.9	11.0	9.1	9.3	10.1	8.6	13.6	15.8	11.9	14.4	16.9	12.5				
1999	9.4	10.3	8.6	8.8	9.5	8.1	13.0	14.8	11.5	13.5	15.5	11.9				
Cerebral infarction (I63)																
2000	5.0	5.1	4.8	4.8	4.9	4.7	5.8	6.5	5.4	6.6	7.2	6.1				
1999	5.5	5.6	5.2	5.3	5.5	5.2	6.2	6.9	5.7	7.1	8.1	6.5				
Stroke, not specified as hemorrhage or infarction (I64)																
2000	32.8	32.2	32.5	31.9	31.1	31.7	38.7	40.6	37.1	44.4	46.8	42.2				
1999	33.7	33.4	33.2	32.8	32.4	32.4	39.3	40.8	37.8	44.9	46.9	42.9				
Other cerebrovascular diseases and their sequelae (I67, I69)																
2000	10.7	11.0	10.4	10.5	10.7	10.2	12.4	12.9	11.9	13.5	13.8	13.1				
1999	10.9	11.2	10.6	10.7	10.9	10.4	12.4	13.5	11.4	13.6	14.5	12.8				

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK	
	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE	MALE	FEMALE
Diseases of arteries, arterioles and capillaries (I70-I78)														
2000.....	14.7	18.1	12.1	14.7	18.3	12.0	13.8	15.6	12.5	15.8	17.8	14.2		
1999.....	15.1	18.3	12.6	15.2	18.5	12.6	13.8	15.6	12.3	15.6	17.7	14.0		
Atherosclerosis (I70)														
2000.....	5.2	5.3	5.0	5.3	5.4	5.1	4.1	4.4	4.0	4.8	5.2	4.6		
1999.....	5.5	5.6	5.4	5.7	5.7	5.5	4.4	4.8	4.1	5.0	5.6	4.6		
Aortic aneurysm and dissection (I71)														
2000.....	5.7	8.6	3.7	5.8	8.9	3.7	4.9	6.3	3.8	5.2	6.6	4.2		
1999.....	5.8	8.7	3.8	5.9	9.0	3.8	4.6	6.1	3.7	5.0	6.3	4.0		
Other diseases of arteries, arterioles and capillaries (I72-I78)														
2000.....	3.7	4.1	3.4	3.6	4.0	3.2	4.8	4.9	4.6	5.7	6.0	5.5		
1999.....	3.7	4.0	3.5	3.6	3.8	3.3	4.7	4.7	4.6	5.6	5.8	5.4		
Other disorders of circulatory system (I80-I99)														
2000.....	1.7	1.7	1.6	1.6	1.6	1.5	2.3	2.3	2.2	2.9	3.0	2.8		
1999.....	1.5	1.5	1.4	1.4	1.4	1.4	2.0	2.2	1.9	2.4	2.6	2.3		
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)														
2000.....	1.2	1.2	1.3	1.2	1.1	1.2	1.7	1.6	1.8	2.2	2.1	2.3		
1999.....	1.0	0.9	1.0	0.9	0.9	0.9	1.3	1.3	1.4	1.7	1.7	1.7		
All other and unspecified disorders of circulatory system (I83-I99)														
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.7	0.4	0.7	0.9	0.5		
1999.....	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.9	0.5	0.7	1.0	0.6		
X. Diseases of the respiratory system (J00-J98)														
2000.....	83.9	103.7	71.6	85.5	104.8	73.7	67.3	90.4	52.7	74.0	99.6	58.2		
1999.....	84.8	106.4	71.5	86.3	107.6	73.5	68.6	92.8	53.3	76.1	103.9	59.3		

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	0.0	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	*	*	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	*	*	*	*
Acute upper respiratory infections (J00-J06)																
2000	0.0	*	*	0.0	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Acute pharyngitis and tonsillitis (J02-J03)																
2000	0.0	*	*	0.0	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Acute laryngitis and tracheitis (J04)																
2000	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	0.0	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)																
2000	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*	*	*	*	*	*	*	*
1999	0.0	0.1	0.0	0.0	0.0	0.0	0.1	*	*	*	0.1	*	*	*	*	*
Other diseases of the respiratory system (J10-J98)																
2000	83.8	103.6	71.6	85.4	104.8	73.6	67.3	90.3	52.7	73.9	99.5	58.2				
1999	84.7	106.3	71.5	86.3	107.5	73.4	68.5	92.7	53.3	76.1	103.7	59.2				
Influenza (J10-J11)																
2000	0.6	0.6	0.6	0.6	0.7	0.7	0.3	0.4	0.3	0.3	0.4	0.3				
1999	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.2	0.3	0.3	0.2	0.3				
Pneumonia (J12-J18)																
2000	23.0	27.5	20.2	22.8	27.0	20.2	23.7	29.9	19.6	25.5	32.3	21.0				
1999	22.9	27.4	20.2	22.8	27.0	20.2	23.0	29.0	19.2	25.3	32.1	21.1				
Viral pneumonia, not elsewhere classified (J12)																
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*			
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK													
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE										
	2000	1.6	2.1	1.3	1.6	2.1	1.2	1.9	2.7	1.5	2.0	2.8	1.5	1999	1.4	1.8	1.2	1.4	1.7	1.2	1.7	2.2	1.3	1.7	2.3	1.4
Bacterial pneumonia (J13-J15)																										
2000	1.6	2.1	1.3	1.6	2.1	1.2	1.9	2.7	1.5	2.0	2.8	1.5		1999	1.4	1.8	1.2	1.4	1.7	1.2	1.7	2.2	1.3	1.7	2.3	1.4
Pneumonia due to other or unspecified organisms (J16,J18)																										
2000	21.3	25.1	18.8	21.1	24.8	18.8	21.6	27.0	18.0	23.3	29.4	19.5		1999	21.4	25.5	18.8	21.2	25.2	18.8	21.2	26.6	17.7	23.4	29.6	19.5
Other acute lower respiratory infections (J20-J22)																										
2000	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.1		1999	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Acute bronchitis and bronchiolitis (J20-J21)																										
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*		1999	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.2	0.2	0.2
Unspecified acute lower respiratory infection (J22)																										
2000	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*	*	*	*		1999	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*	*	*	*
Other diseases of upper respiratory tract (J30-J39)																										
2000	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.3		1999	0.2	0.2	0.1	0.1	0.2	0.3	0.2	0.3	0.3	0.3	0.2	0.2
Chronic lower respiratory diseases (J40-J47)																										
2000	44.3	54.9	37.9	46.1	56.3	40.0	28.6	41.5	20.6	31.7	46.0	23.1		1999	45.8	58.1	38.2	47.5	59.6	40.2	30.3	44.6	21.4	33.7	50.1	23.9
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)																										
2000	44.0	54.7	37.4	45.7	56.1	39.5	28.4	41.3	20.4	31.6	45.9	23.0		1999	45.4	57.8	37.8	47.1	59.3	39.8	30.1	44.3	21.2	33.6	50.0	23.7

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Bronchitis, not specified as acute or chronic (J40)															
2000.....	0.2	0.2	0.2	0.2	0.3	0.2	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2
Chronic bronchitis (J41-J42)															
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	*	0.1	*	*	*	*	*
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	*	0.1	*	*	*	*	*
Emphysema (J43)															
2000.....	6.1	7.7	5.0	6.4	8.0	5.4	3.4	5.5	2.1	3.8	6.0	2.4			
1999.....	6.5	8.6	5.2	6.9	8.9	5.7	3.6	5.9	2.1	4.0	6.6	2.4			
Other chronic obstructive pulmonary disease (J44)															
2000.....	35.8	45.1	30.2	37.6	46.6	32.2	21.3	32.4	14.5	23.6	36.0	16.1			
1999.....	36.7	47.4	30.2	38.4	48.9	32.1	22.7	34.7	15.2	25.2	39.4	16.8			
Asthma (J45-J46)															
2000.....	1.6	1.3	1.8	1.3	1.0	1.5	3.3	3.1	3.5	3.9	3.5	4.2			
1999.....	1.7	1.4	2.0	1.4	1.0	1.7	3.4	3.3	3.5	3.9	3.6	4.2			
Bronchiectasis (J47)															
2000.....	0.3	0.2	0.4	0.4	0.2	0.4	0.2	0.2	0.3	0.1	*	*			
1999.....	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.2	*	0.2			
Lung diseases due to external agents (J60-J70)															
2000.....	6.5	9.4	4.7	6.5	9.5	4.7	6.2	8.5	4.7	7.1	9.8	5.4			
1999.....	6.1	9.0	4.4	6.1	9.0	4.4	6.0	8.8	4.4	7.2	10.3	5.3			
Pneumoconioses and chemical effects (J60-J66,J68)															
2000.....	0.4	1.1	0.0	0.4	1.1	0.0	0.2	0.5	*	0.3	0.6	*			
1999.....	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.6	*	0.3	0.7	*			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK												
	BOTH SEXES		MALE																						
	2000	6.0	8.3	4.7	6.0	8.4	4.7	5.9	7.9	4.7	6.8	9.1	5.4	1999	5.6	7.9	4.4	5.6	7.9	4.3	5.8	8.1	4.4	6.8	9.5
Pneumonitis due to solids and liquids (J69)																									
2000	6.0	8.3	4.7	6.0	8.4	4.7	5.9	7.9	4.7	6.8	9.1	5.4													
1999	5.6	7.9	4.4	5.6	7.9	4.3	5.8	8.1	4.4	6.8	9.5	5.2													
Pneumonitis due to food and vomit (J69.0)																									
2000	6.0	8.3	4.7	6.0	8.4	4.7	5.9	7.9	4.7	6.8	9.1	5.4													
1999	5.6	7.9	4.4	5.6	7.9	4.3	5.8	8.1	4.4	6.8	9.5	5.2													
Other pneumonitis due to solids and liquids (J69.1-J69.8)																									
2000	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*										
1999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*										
Other lung diseases due to external agents (J67,J70)																									
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*										
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*										
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)																									
2000	0.2	0.4	0.2	0.2	0.3	0.2	0.4	0.6	0.3	0.5	0.7	0.3													
1999	0.3	0.4	0.2	0.3	0.4	0.2	0.4	0.7	0.3	0.5	0.8	0.3													
Pleural effusion and plaque (J90,J92)																									
2000	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.2	0.4	0.5	0.3													
1999	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4													
All other diseases of respiratory system (J80-J84,J93-J98)																									
2000	8.4	10.0	7.4	8.5	10.2	7.4	7.5	8.8	6.6	8.2	9.4	7.4													
1999	8.3	10.0	7.3	8.4	10.1	7.2	7.6	8.6	7.0	8.3	9.3	7.6													
XI. Diseases of the digestive system (K00-K92)																									
2000	30.5	35.6	25.8	30.2	35.1	25.5	31.5	38.1	26.2	35.5	43.5	29.3													
1999	30.6	36.0	25.7	30.2	35.5	25.4	31.8	38.7	26.3	35.7	44.0	29.3													

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Diseases of oral cavity, salivary glands and jaws (K00-K14)															
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*
Diseases of other parts of digestive system (K20-K92)															
2000.....	30.6	35.8	25.8	30.3	35.3	25.5	31.5	38.3	26.2	35.6	43.6	29.3			
1999.....	30.7	36.3	25.6	30.4	35.8	25.4	31.8	38.9	26.2	35.8	44.2	29.2			
Diseases of esophagus (K20-K22)															
2000.....	0.8	1.0	0.7	0.8	1.0	0.7	0.6	0.8	0.4	0.7	1.0	0.5			
1999.....	0.8	1.0	0.7	0.8	1.0	0.7	0.6	0.8	0.4	0.7	1.0	0.5			
Peptic ulcer (K25-K28)															
2000.....	1.6	2.0	1.4	1.6	2.0	1.4	1.7	2.3	1.2	1.7	2.3	1.4			
1999.....	1.7	2.0	1.4	1.7	2.0	1.4	1.8	2.2	1.4	1.9	2.4	1.4			
Gastric ulcer (K25)															
2000.....	0.5	0.5	0.4	0.5	0.5	0.4	0.6	0.9	0.4	0.7	1.0	0.4			
1999.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.9	0.5	0.7	1.0	0.5			
Duodenal ulcer (K26)															
2000.....	0.5	0.7	0.4	0.6	0.7	0.4	0.5	0.6	0.4	0.5	0.6	0.4			
1999.....	0.6	0.7	0.4	0.5	0.7	0.4	0.5	0.6	0.4	0.5	0.6	0.4			
Peptic ulcer, site unspecified (K27)															
2000.....	0.6	0.8	0.5	0.6	0.8	0.5	0.5	0.7	0.4	0.6	0.6	0.5			
1999.....	0.7	0.8	0.6	0.6	0.7	0.6	0.6	0.7	0.5	0.6	0.8	0.5			
Gastrojejunal ulcer (K28)															
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*			
1999.....	0.0	*	*	0.0	*	*	*	*	*	*	*	*			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Gastritis and duodenitis (K29)															
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	*	0.1		
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1	0.2	0.3	0.3	0.2		
Dyspepsia and other diseases of stomach and duodenum (K30-K31)															
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2		
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2		
Diseases of appendix (K35-K38)															
2000.....	0.2	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.1	0.3	0.4	0.4	0.2		
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1	0.2	0.4	0.4	0.1		
Hernia (K40-K46)															
2000.....	0.6	0.5	0.6	0.6	0.5	0.6	0.5	0.5	0.5	0.6	0.6	0.6	0.6		
1999.....	0.5	0.5	0.5	0.5	0.6	0.5	0.5	0.5	0.5	0.6	0.6	0.6	0.6		
Crohn's disease and ulcerative colitis (K50-K51)															
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.1	0.2	0.2	*	0.2			
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.1	0.2	0.2	0.2	0.2			
Vascular disorders and obstruction of intestine without hernia (K55-K56)															
2000.....	5.2	4.6	5.5	5.1	4.4	5.5	5.6	5.9	5.3	6.6	7.1	6.2			
1999.....	5.2	4.5	5.5	5.1	4.4	5.5	5.4	5.5	5.4	6.5	6.5	6.4			
Diverticular disease of intestine (K57)															
2000.....	1.2	1.0	1.3	1.2	1.0	1.4	0.9	0.6	1.0	1.0	0.8	1.2			
1999.....	1.3	1.0	1.4	1.3	1.0	1.4	0.9	0.7	1.0	1.1	0.8	1.3			
Other diseases of intestines and peritoneum (K52,K58-K63)															
2000.....	1.6	1.5	1.7	1.6	1.5	1.7	1.6	1.7	1.5	1.8	1.8	1.7			
1999.....	1.6	1.6	1.7	1.6	1.5	1.7	1.5	1.7	1.4	1.7	1.9	1.6			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER		
	BOTH SEXES		MALE									
	BOTH SEXES	MALE	FEMALE									
Diseases of peritoneum (K65-K66)												
2000.....	0.7	0.7	0.7	0.7	0.7	0.6	0.9	1.1	0.9	1.2	1.3	1.1
1999.....	0.7	0.7	0.6	0.7	0.7	0.6	1.0	1.0	0.9	1.1	1.2	1.0
Diseases of liver (K70-K76)												
2000.....	12.3	16.8	8.3	12.2	16.8	8.1	12.3	16.6	8.9	13.3	18.6	9.2
1999.....	12.4	17.2	8.2	12.3	17.0	8.0	13.0	17.7	9.1	14.2	19.9	9.7
Alcoholic liver disease (K70)												
2000.....	4.4	6.9	2.2	4.4	6.9	2.1	4.4	6.6	2.7	4.7	7.1	2.8
1999.....	4.4	7.0	2.1	4.3	6.9	2.0	4.8	7.5	2.7	5.2	8.2	2.8
Chronic hepatitis, not elsewhere classified (K73)												
2000.....	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.2	0.2	*	0.2
1999.....	0.2	0.1	0.2	0.1	0.1	0.2	0.2	0.1	0.3	0.2	*	0.3
Fibrosis and cirrhosis of liver (K74)												
2000.....	5.1	6.5	3.8	5.2	6.6	3.9	4.4	5.8	3.3	4.7	6.5	3.3
1999.....	5.1	6.6	3.8	5.2	6.7	3.9	4.5	5.9	3.2	4.8	6.8	3.3
Other diseases of liver (K71-K72, K75-K76)												
2000.....	2.7	3.3	2.1	2.5	3.2	1.9	3.3	4.2	2.6	3.8	4.9	2.9
1999.....	2.7	3.5	2.1	2.6	3.3	1.9	3.5	4.1	2.9	3.9	4.8	3.2
Cholelithiasis and other disorders of gallbladder (K80-K82)												
2000.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.0	1.1	1.2	1.0	1.3
1999.....	1.0	1.2	1.0	1.0	1.2	0.9	1.1	1.2	1.1	1.1	1.1	1.1
Disorders of biliary tract and pancreas (K83-K86)												
2000.....	1.7	2.0	1.4	1.6	1.9	1.3	2.1	2.8	1.7	2.5	3.3	1.9
1999.....	1.6	2.0	1.3	1.5	1.9	1.3	2.1	2.7	1.6	2.4	3.2	1.8

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Pancreatitis (K85,K86.0-K86.1)															
2000.....	1.2	1.5	1.0	1.2	1.4	0.9	1.6	2.2	1.2	1.9	2.6	1.4			
1999.....	1.2	1.5	1.0	1.1	1.4	0.9	1.6	2.1	1.2	1.9	2.6	1.3			
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)															
2000.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.6	0.5	0.6	0.7	0.6			
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.6	0.4	0.5	0.6	0.5			
All other diseases of digestive system (K90,K92)															
2000.....	2.9	3.3	2.6	2.9	3.2	2.5	3.4	3.9	3.0	4.0	4.7	3.5			
1999.....	2.9	3.3	2.5	2.8	3.2	2.5	3.1	3.7	2.7	3.6	4.2	3.1			
XII. Diseases of the skin and subcutaneous tissue (L00-L98)															
2000.....	1.3	1.2	1.5	1.1	1.0	1.2	3.2	2.8	3.4	3.9	3.6	4.1			
1999.....	1.3	1.2	1.4	1.1	1.0	1.2	3.4	3.0	3.6	4.3	3.8	4.5			
Infections of skin and subcutaneous tissue (L00-L08)															
2000.....	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5	0.5	0.5	0.5			
1999.....	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.4	0.5	0.6	0.5			
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)															
2000.....	0.9	0.8	1.1	0.7	0.6	0.8	2.7	2.3	2.9	3.4	3.0	3.5			
1999.....	1.0	0.9	1.1	0.8	0.7	0.8	3.0	2.5	3.3	3.8	3.2	4.0			
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)															
2000.....	5.0	3.6	6.0	4.8	3.5	5.7	5.7	4.6	6.6	6.6	5.6	7.6			
1999.....	4.9	3.5	5.9	4.7	3.4	5.7	5.5	4.1	6.6	6.3	4.9	7.5			
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)															
2000.....	1.0	0.6	1.3	1.0	0.6	1.3	0.6	0.3	0.8	0.6	0.4	0.8			
1999.....	1.0	0.6	1.3	1.0	0.6	1.4	0.6	0.3	0.7	0.6	0.4	0.7			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK												
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE									
	2000	0.5	0.2	0.8	0.3	0.1	0.6	1.1	0.3	1.8	1.4	0.4	2.2	1999	0.5	0.2	0.8	0.4	0.1	0.6	1.1	0.3	1.8	1.3	0.3
Systemic lupus erythematosus (M32)																									
2000	0.5	0.2	0.8	0.3	0.1	0.6	1.1	0.3	1.8	1.4	0.4	2.2													
1999	0.5	0.2	0.8	0.4	0.1	0.6	1.1	0.3	1.8	1.3	0.3	2.2													
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)																									
2000	1.7	1.3	2.1	1.7	1.2	2.0	2.1	1.8	2.3	2.4	2.1	2.7													
1999	1.7	1.3	2.0	1.6	1.2	1.9	2.1	1.6	2.5	2.5	1.9	2.9													
Dorsopathies (M40-M54)																									
2000	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.2													
1999	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.2													
Soft tissue disorders (M60-M79)																									
2000	0.4	0.5	0.3	0.3	0.4	0.2	0.7	0.7	0.6	0.8	0.9	0.7													
1999	0.4	0.5	0.3	0.3	0.4	0.2	0.6	0.8	0.5	0.7	1.0	0.5													
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)																									
2000	1.1	0.8	1.2	1.1	0.8	1.3	1.0	1.1	0.9	1.1	1.3	0.9													
1999	1.1	0.7	1.2	1.1	0.7	1.3	0.8	0.8	0.8	0.9	0.9	0.9													
Osteoporosis (M80-M81)																									
2000	0.6	0.2	0.8	0.7	0.2	0.9	0.2	*	0.2	0.1	*	*													
1999	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	0.2	0.1	*	*													
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)																									
2000	0.5	0.6	0.4	0.4	0.5	0.4	0.8	1.0	0.7	1.0	1.2	0.8													
1999	0.4	0.5	0.4	0.4	0.5	0.4	0.7	0.7	0.6	0.8	0.9	0.8													
XIV. Diseases of the genitourinary system (N00-N98)																									
2000	19.8	22.5	18.1	18.1	20.9	16.5	31.5	33.7	30.0	37.9	41.3	35.6													
1999	19.6	22.4	17.9	17.9	20.6	16.3	32.1	35.6	29.7	38.2	43.3	35.0													

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK													
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE										
	2000	19.5	21.9	17.9	17.8	20.4	16.3	31.1	33.0	29.6	37.4	40.5	35.2	1999	19.3	21.7	17.7	17.6	20.0	16.1	31.6	35.0	29.3	37.6	42.5	34.5
Diseases of urinary system (N00-N39)																										
2000	0.7	0.8	0.7	0.7	0.7	0.6	1.0	1.1	1.0	1.0	1.1	1.2	1.0	1999	0.8	0.9	0.7	0.7	0.8	0.7	1.0	1.2	0.9	1.1	1.4	0.9
Glomerular and renal tubulo-interstitial diseases (N00-N15)																										
2000	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.4	0.4	0.4	0.5	0.4	1999	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4	0.5	0.6	0.4
Glomerular diseases (N00-N07)																										
2000	0.0	*	*	0.0	*	*	*	*	*	*	*	*	*	1999	0.0	*	*	0.0	*	*	*	*	*	*	*	*
Acute and rapidly progressive nephritic syndrome (N00-N01)																										
2000	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3	0.3	0.4	0.4	0.3	1999	0.2	0.3	0.1	0.2	0.2	0.1	0.4	0.4	0.3	0.4	0.4	0.3
Nephrotic syndrome (N04)																										
2000	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*	0.1	*	*	*	1999	0.1	0.1	0.0	0.0	0.0	0.1	*	*	0.1	*	*	*
Other and unspecified glomerular diseases (N02-N03,N05-N07)																										
2000	0.5	0.5	0.5	0.4	0.4	0.4	0.6	0.6	0.6	0.6	0.7	0.7	0.6	1999	0.5	0.6	0.5	0.5	0.5	0.5	0.8	0.8	0.6	0.7	0.6	0.3
Renal tubulo-interstitial diseases (N10-N15)																										
2000	0.5	0.5	0.5	0.4	0.4	0.4	0.6	0.6	0.6	0.6	0.7	0.7	0.6	1999	0.5	0.6	0.5	0.5	0.5	0.5	0.8	0.8	0.6	0.7	0.6	0.6
Renal failure (N17-N19)																										
2000	13.2	16.3	11.4	11.8	14.9	9.9	23.2	25.5	21.6	28.3	31.7	26.1	25.5	1999	12.8	15.9	11.0	11.4	14.5	9.5	23.4	26.4	21.5	28.1	32.3	25.5

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	0.2	...	0.3	0.1	...	0.3
1999	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	0.3	...	0.4	0.1	...	0.4	
Disorders of female genital organs (N70-N98)																
2000	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	0.1	...	0.1	0.1	...	0.1	
1999	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
Inflammatory diseases of female pelvic organs (N70-N76)																
2000	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	
1999	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	0.2	...	0.2	0.2	...	0.3	
Noninflammatory disorders of female genital tract (N80-N98)																
2000	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	
1999	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	0.2	...	0.2	0.2	...	0.3	
XV. Pregnancy, childbirth and the puerperium (000-099)																
2000	0.1	...	0.3	0.1	...	0.2	0.3	...	0.6	0.4	...	0.7	0.4	...	0.8	
1999	0.2	...	0.3	0.1	...	0.2	0.4	...	0.7	0.4	...	0.8	0.4	...	0.8	
Pregnancy with abortive outcome (000-007)																
2000	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*	
1999	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*	
Ectopic pregnancy (000)																
2000	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*	
1999	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*	
Spontaneous abortion (003)																
2000	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*	
1999	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*	
Medical abortion (004)																
2000	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*	
1999	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*	

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Other abortion (005)															
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	...	*	
Other and unspecified pregnancy with abortive outcome (001-002,006-007)															
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	...	*	
Other direct obstetric deaths (010-092)															
2000.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.4	0.3	...	0.5	...		
1999.....	0.1	...	0.2	0.1	...	0.1	0.3	...	0.5	0.3	...	0.6	...		
Eclampsia and pre-eclampsia (011,013-016)															
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	0.1	...	0.1	...	0.1	
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2	0.1	...	0.2	...	0.2	
Hemorrhage of pregnancy and childbirth and placenta previa (020,044-046,067,072)															
2000.....	0.0	...	0.0	0.0	...	0.0	*	...	*	*	...	*	...	*	
1999.....	0.0	...	0.0	0.0	...	0.0	*	...	*	*	...	*	...	*	
Complications predominantly related to the puerperium (085-092)															
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	0.1	...	0.1	...	0.1	
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1	0.1	...	0.1	...	0.2	
Obstetric embolism (088)															
2000.....	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1	*	...	*	...	*	
1999.....	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1	*	...	*	...	*	
Other complications predominately related to the puerperium (085-087,089-092)															
2000.....	0.0	...	0.0	*	...	*	*	...	*	*	...	*	...	*	
1999.....	0.0	...	0.0	*	...	*	*	...	*	*	...	*	...	*	

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.4	0.4	0.4	0.4	0.4
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)	1999	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.4	0.4	0.4
Congenital malformations of heart (Q20-Q24)																
2000	1.2	1.3	1.1	1.2	1.3	1.1	1.3	1.4	1.2	1.5	1.6	1.4	1.5	1.6	1.4	1.4
1999	1.3	1.4	1.2	1.3	1.4	1.2	1.3	1.5	1.2	1.6	1.7	1.6	1.7	1.4	1.4	1.4
Other congenital malformations of circulatory system (Q25-Q28)																
2000	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.5	0.4	0.5	0.4	0.4	0.4
1999	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.5	0.5	0.5	0.5	0.4	0.4	0.4
Congenital malformations of respiratory system (Q30-Q34)																
2000	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.4
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.4	0.4	0.4	0.4	0.4	0.2
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)																
2000	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.1	0.8	1.1	1.3	1.0	1.1	1.3	1.0	1.0
1999	0.9	1.1	0.8	0.9	1.0	0.8	1.0	1.1	0.9	1.1	1.3	1.1	1.3	1.0	1.0	1.0
Down's syndrome (Q90)																
2000	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)																
2000	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.5	0.4	0.5	0.4	0.5	0.5	0.5
1999	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5	0.4	0.3	0.4	0.3	0.4	0.3	0.5
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)																
2000	11.5	13.1	9.8	11.0	12.5	9.3	13.7	16.0	11.5	16.8	19.8	14.0				
1999	9.8	10.9	8.5	9.1	10.1	8.0	12.8	15.3	10.6	15.7	19.2	12.8				

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK													
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE										
	2000	1.4	1.1	1.5	1.4	1.1	1.5	1.2	0.8	1.3	1.4	1.0	1.5	1999	1.2	1.0	1.2	1.2	1.0	1.3	0.9	0.8	0.9	1.0	1.0	1.0
Senility (R54)																										
2000	1.4	1.1	1.5	1.4	1.1	1.5	1.2	0.8	1.3	1.4	1.0	1.5		1999	1.2	1.0	1.2	1.2	1.0	1.3	0.9	0.8	0.9	1.0	1.0	1.0
Ill-defined and unknown causes of mortality (R95-R99)																										
2000	7.4	9.2	5.8	7.0	8.7	5.4	9.0	11.4	7.1	11.1	14.1	8.7		1999	6.1	7.3	4.8	5.5	6.6	4.4	8.5	10.9	6.5	10.4	13.5	7.9
Sudden infant death syndrome (R95)																										
2000	0.9	1.1	0.7	0.7	0.9	0.6	1.6	1.8	1.3	2.0	2.2	1.7		1999	1.0	1.1	0.8	0.8	0.9	0.7	1.6	1.9	1.4	2.0	2.3	1.8
Other ill-defined and unknown causes of mortality (R96-R99)																										
2000	6.5	8.1	5.0	6.3	7.9	4.8	7.5	9.6	5.8	9.2	11.9	7.0		1999	5.1	6.2	4.0	4.7	5.7	3.8	6.8	9.0	5.1	8.4	11.2	6.1
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)																										
2000	2.7	2.8	2.5	2.6	2.6	2.4	3.5	3.8	3.1	4.3	4.8	3.8		1999	2.6	2.6	2.5	2.4	2.5	2.3	3.5	3.6	3.2	4.3	4.7	3.9
XX. External causes of mortality (V01-Y89)																										
2000	54.6	80.5	31.0	53.1	77.3	30.7	60.2	93.8	31.0	69.2	110.4	33.8		1999	55.2	81.3	31.5	53.4	77.8	31.1	62.2	96.5	32.7	71.2	113.2	35.5
Accidents (unintentional injuries) (V01-X59,Y85-Y86)																										
2000	35.5	50.0	22.3	35.5	49.7	22.5	34.7	51.6	20.6	38.4	58.8	21.8		1999	35.9	50.6	22.7	35.7	50.0	22.7	36.6	54.2	22.0	40.9	62.1	23.7
Transport accidents (V01-V99,Y85)																										
2000	16.9	24.2	10.2	17.1	24.3	10.4	16.1	24.2	9.2	17.0	26.6	9.0		1999	17.0	24.1	10.4	17.1	24.1	10.5	16.6	24.5	9.9	17.6	27.0	9.7

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK												
	BOTH SEXES		MALE																						
	2000	0.2	0.3	0.1	0.2	0.3	0.0	0.2	0.4	*	0.3	0.5	*	1999	0.2	0.3	0.0	0.2	0.3	0.1	0.2	0.4	*	0.2	0.4
Railway accidents (V05,V15,V80.6,V81.2-V81.9)																									
2000.....	0.2	0.3	0.1	0.2	0.3	0.0	0.2	0.4	*	0.3	0.5	*													
1999.....	0.2	0.3	0.0	0.2	0.3	0.1	0.2	0.4	*	0.2	0.4	*													
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)																									
2000.....	15.7	22.2	9.7	15.9	22.2	9.9	15.1	22.5	8.9	16.0	24.8	8.6													
1999.....	15.5	21.8	9.8	15.6	21.8	9.9	15.3	22.3	9.3	16.2	24.7	9.2													
Pedestrian involved in collision with motor vehicle (V02-V04)																									
2000.....	1.2	1.8	0.8	1.1	1.6	0.7	1.8	2.8	0.9	1.8	3.0	0.9													
1999.....	1.3	1.8	0.7	1.1	1.6	0.7	1.9	3.0	1.0	2.0	3.2	1.0													
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)																									
2000.....	0.2	0.4	0.0	0.2	0.4	0.0	0.3	0.6	*	0.3	0.7	*													
1999.....	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	*	0.3	0.5	*													
Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)																									
2000.....	1.0	1.8	0.2	1.1	1.9	0.2	0.7	1.5	*	0.9	1.8	*													
1999.....	0.9	1.6	0.2	0.9	1.6	0.2	0.6	1.2	*	0.7	1.5	*													
Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)																									
2000.....	0.1	0.1	0.0	0.1	0.2	0.0	0.1	0.1	*	0.1	*	*													
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.2	*													
Motorcyclist involved in collision with railway train (V25,V35)																									
2000.....	*	*	*	*	*	*	*	*	*	*	*	*													
1999.....	*	*	*	*	*	*	*	*	*	*	*	*													
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)																									
2000.....	0.1	0.1	0.0	0.1	0.2	0.0	0.1	0.1	*	0.1	*	*													
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	*													

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES		MALE	BOTH SEXES		MALE									
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)															
2000.....	1.4	2.1	0.8	1.4	2.0	0.8	1.5	2.4	0.7	1.6	2.7	0.7			
1999.....	1.4	2.1	0.7	1.4	2.1	0.7	1.5	2.3	0.8	1.7	2.7	0.9			
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)															
2000.....	3.8	4.7	2.9	3.8	4.8	3.0	3.4	4.5	2.4	3.5	4.9	2.4			
1999.....	3.8	4.7	2.9	3.8	4.7	3.0	3.4	4.7	2.5	3.7	5.2	2.5			
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)															
2000.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	*	0.1	0.2	*			*
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	0.1	*	*			*
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)															
2000.....	1.4	2.0	0.8	1.4	2.0	0.8	1.2	1.8	0.8	1.1	1.8	0.6			
1999.....	1.4	1.9	0.8	1.4	1.9	0.8	1.3	1.8	0.8	1.2	1.8	0.7			
Occupant of special-use motor vehicle involved in any accident (V83-V86)															
2000.....	0.4	0.7	0.1	0.4	0.8	0.1	0.2	0.3	*	0.1	0.3	*			*
1999.....	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.3	*	0.1	0.3	*			*
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)															
2000.....	6.2	8.4	4.1	6.2	8.4	4.2	6.0	8.4	3.9	6.3	9.4	3.8			
1999.....	6.1	8.3	4.2	6.2	8.3	4.3	5.9	8.3	4.0	6.4	9.3	4.0			
Streetcar accidents (V82.2-V82.9)															
2000.....	*	*	*	*	*	*	*	*	*	*	*	*			*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*			*
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)															
2000.....	0.4	0.5	0.2	0.4	0.5	0.2	0.3	0.5	0.2	0.3	0.5	0.2			0.2
1999.....	0.6	0.9	0.3	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.9	0.3			0.3

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK												
	BOTH SEXES		MALE	BOTH SEXES		MALE																			
	2000	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*	0.2	0.4	*	1999	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.5	*	0.3	0.5
Water transport accidents (V90-V94)																									
2000	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*	0.2	0.4	*													
1999	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.5	*	0.3	0.5	*													
Air and space transport accidents (V95-V97)																									
2000	0.3	0.5	0.1	0.3	0.5	0.1	0.1	0.1	*	*	*	*													
1999	0.3	0.4	0.1	0.3	0.5	0.1	0.0	0.1	*	*	*	*													
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)																									
2000	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*	0.2	0.4	*													
1999	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1	0.3	0.4	0.1													
Falls (W00-W19)																									
2000	4.8	6.5	3.6	5.0	6.7	3.8	3.1	4.6	2.1	3.0	4.5	1.9													
1999	4.8	6.6	3.6	5.0	6.7	3.7	3.5	5.3	2.2	3.5	5.2	2.2													
Fall on same level (W00-W09,W18)																									
2000	1.2	1.4	1.0	1.2	1.5	1.0	0.7	0.9	0.5	0.5	0.7	0.4													
1999	0.8	1.0	0.7	0.8	1.0	0.7	0.6	0.8	0.5	0.6	0.8	0.5													
Fall from one level to another (W10-W17)																									
2000	1.1	1.8	0.5	1.1	1.9	0.5	0.7	1.3	0.3	0.7	1.3	0.3													
1999	1.1	1.9	0.5	1.2	2.0	0.6	0.8	1.4	0.3	0.8	1.3	0.3													
Unspecified fall (W19)																									
2000	2.6	3.3	2.1	2.7	3.3	2.2	1.7	2.5	1.3	1.7	2.5	1.2													
1999	2.9	3.7	2.4	3.0	3.7	2.4	2.0	3.0	1.4	2.1	3.1	1.4													
Accidental discharge of firearms (W32-W34)																									
2000	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	0.1	0.3	0.7	0.1													
1999	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	0.1	0.4	0.8	0.1													

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK			
	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE	BOTH SEXES		MALE	FEMALE
	2000	0.1	0.1	0.0	0.1	0.1	0.0	0.0	*	*	*	*	*	*	*	*
Accidental explosion (W39-W40)	1999	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*	*	*	*	*	*	*
Accidental drowning and submersion (W65-W74)																
2000	1.3	2.0	0.5	1.2	1.9	0.5	1.6	2.6	0.7	1.6	2.7	0.6				
1999	1.3	2.1	0.5	1.2	1.9	0.5	1.6	2.8	0.6	1.7	3.0	0.6				
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)																
2000	1.6	2.0	1.3	1.5	1.9	1.3	1.8	2.4	1.4	2.2	2.9	1.7				
1999	1.4	1.8	1.2	1.4	1.7	1.2	1.8	2.2	1.5	2.2	2.8	1.8				
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)																
2000	0.5	0.7	0.3	0.5	0.7	0.3	0.5	0.7	0.4	0.6	0.8	0.5				
1999	0.6	0.8	0.4	0.6	0.8	0.4	0.7	0.9	0.5	0.8	1.0	0.5				
Accidental exposure to electric current (W85-W87)																
2000	0.1	0.3	0.0	0.2	0.3	*	0.1	0.1	*	0.1	0.2	*	0.1	0.2	*	*
1999	0.2	0.3	0.0	0.2	0.3	0.0	0.1	0.2	*	0.1	0.2	*	0.1	0.2	*	*
Accidental exposure to radiation (W88-W91)																
2000	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)																
2000	1.2	1.6	0.9	1.0	1.3	0.8	2.4	3.2	1.8	3.1	4.2	2.3				
1999	1.2	1.6	0.9	1.0	1.4	0.7	2.4	3.2	1.9	3.0	4.0	2.3				
Accidental contact with heat and hot substances (X10-X19)																
2000	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	0.1	*	*	0.1	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	*	0.1	*	0.1	*	*	*	*

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE												
Lightning (X33)															
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.1	*	0.0	0.1	*	*	*	*	*	*	*	*	*	*
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)															
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*	*	*	*	*	*	*
Cataclysmic storm and flood (X37-X38)															
2000.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*	0.1	*	*	*	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)															
2000.....	4.6	6.7	2.6	4.6	6.7	2.5	4.8	7.3	2.7	6.2	9.5	3.4			
1999.....	4.5	6.6	2.3	4.4	6.5	2.3	5.1	7.8	2.7	6.4	10.1	3.3			
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)															
2000.....	4.2	6.1	2.4	4.2	6.1	2.3	4.4	6.7	2.5	5.7	8.8	3.1			
1999.....	4.1	6.0	2.2	4.0	5.9	2.1	4.6	7.1	2.4	5.9	9.3	3.0			
Accidental poisoning by and exposure to other and unspecified solid or liquid substances (X45-X46,X48-X49)															
2000.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	0.1	0.2	0.3	*			*
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	0.1	0.2	0.3	*			*
Accidental poisoning by and exposure to other gases and vapors (X47)															
2000.....	0.2	0.4	0.1	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1			
1999.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1			
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)															
2000.....	4.0	5.3	2.8	4.0	5.3	2.8	3.6	5.6	2.1	4.1	6.5	2.3			
1999.....	4.4	5.9	3.1	4.4	5.8	3.2	4.3	6.4	2.7	5.0	7.6	3.1			

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR				ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK		
	BOTH SEXES	MALE	FEMALE															
Intentional self-harm (suicide) (X60-X84,Y87.0)																		
2000.....	10.6	18.1	4.0	11.5	19.4	4.4	6.0	10.4	2.2	5.6	10.2	1.8						
1999.....	10.7	18.2	4.1	11.5	19.4	4.4	6.2	10.7	2.2	5.7	10.4	1.6						
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)																		
2000.....	1.2	1.1	1.2	1.3	1.3	1.4	0.5	0.5	0.4	0.5	0.5	0.4						
1999.....	1.2	1.1	1.2	1.3	1.3	1.3	0.5	0.5	0.4	0.4	0.5	0.4						
Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)																		
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	0.0	*	*	*	*	*						
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	0.0	*	*	*	*	*						
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)																		
2000.....	0.5	0.9	0.2	0.6	1.0	0.3	0.1	0.2	*	0.1	0.2	*					*	
1999.....	0.6	0.9	0.3	0.6	1.1	0.3	0.2	0.2	*	0.1	0.2	*				*		
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)																		
2000.....	2.1	3.5	0.7	2.2	3.7	0.7	1.6	2.8	0.6	1.2	2.1	0.3						
1999.....	2.0	3.4	0.7	2.1	3.5	0.7	1.6	2.7	0.6	1.0	2.0	0.2						
Intentional self-harm (suicide) by discharge of firearms (X72-X74)																		
2000.....	6.0	11.2	1.5	6.6	12.1	1.7	2.9	5.5	0.7	3.2	6.1	0.7						
1999.....	6.1	11.4	1.5	6.6	12.2	1.7	3.1	6.0	0.6	3.3	6.6	0.6						
Intentional self-harm (suicide) by jumping from a high place (X80)																		
2000.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	0.1					*	
1999.....	0.3	0.4	0.1	0.3	0.4	0.1	0.3	0.4	0.2	0.2	0.3	0.3				*		
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (X71,X75-X79,X81-X84,Y87.0)																		
2000.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.9	0.3	0.5	0.8	0.2						
1999.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.3	0.5	0.8	0.3						

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992) AND YEAR	ALL RACES			WHITE			TOTAL			ALL OTHER			BLACK													
	BOTH SEXES		MALE	BOTH SEXES		MALE																				
	2000	0.4	0.5	0.2	0.3	0.4	0.2	0.6	0.9	0.3	0.6	1.0	0.3	1999	0.3	0.5	0.2	0.3	0.5	0.2	0.8	0.2	0.6	0.6	0.9	0.2
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21, Y25-Y29, Y31-Y34, Y87.2, Y89.9)																										
2000	0.4	0.5	0.2	0.3	0.4	0.2	0.6	0.9	0.3	0.6	1.0	0.3	*	1999	0.3	0.5	0.2	0.3	0.5	0.2	0.8	0.2	0.6	0.6	0.9	0.2
Legal intervention (Y35, Y89.0)																										
2000	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*	0.3	0.6	*	*	1999	0.2	0.3	*	0.1	0.2	*	0.3	0.7	*	*	*	*
Legal execution (Y35.5)																										
2000	0.0	0.0	*	0.0	0.0	*	0.1	0.1	*	0.1	0.2	*	*	1999	0.0	0.1	*	0.0	0.1	*	0.1	0.2	*	*	*	*
Other legal intervention and their sequelae (Y35.0-Y35.4, Y35.6-Y35.7, Y89.0)																										
2000	0.1	0.2	*	0.1	0.2	*	0.2	0.3	*	0.2	0.4	*	*	1999	0.1	0.2	*	0.1	0.2	*	0.3	0.5	*	*	*	*
Operations of war and their sequelae (Y36, Y89.1)																										
2000	*	*	*	*	*	*	*	*	*	*	*	*	*	1999	0.0	0.0	*	0.0	0.0	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84, Y88)																										
2000	1.1	1.2	1.0	1.1	1.2	0.9	1.5	1.5	1.6	1.9	1.8	2.0	*	1999	1.0	1.2	1.0	1.0	1.1	0.9	1.5	1.7	1.4	1.8	1.9	1.7
Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59, Y88.0)																										
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	*	*	1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.2	*	*
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69, Y88.1)																										
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	*	*	1999	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	*	0.2	0.2

02/10/24 WORK TABLE 13. AGE-ADJUSTED DEATH RATES FOR 358 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1999-2000 PAGE 58

(AGE-ADJUSTED DEATH RATES PER 100,000 U.S. STANDARD POPULATION BASED ON YEAR 2000 STANDARD IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION,
INTERNATIONAL CLASSIFICATION OF DISEASES, 1992)
- - ALL RACES - - - - - WHITE - - - - - TOTAL - - - - - ALL OTHER - - - - - BLACK - - - - -
AND YEAR BOTH SEXES MALE FEMALE BOTH SEXES MALE FEMALE BOTH SEXES MALE FEMALE BOTH SEXES MALE FEMALE

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

2000.....	1.0	1.1	0.9	0.9	1.0	0.8	1.3	1.3	1.3	1.6	1.6	1.6
1999.....	0.9	1.0	0.8	0.8	0.9	0.8	1.3	1.5	1.2	1.5	1.7	1.4

* FIGURE DOES NOT MEET STANDARDS OF RELIABILITY OR PRECISION.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.